

Dr. Rishma Dhillon Pai uses MRI guided focused ultrasound at Jaslok Hospital to treat fibroids without surgery, says Aditi Mallya

libeeds are common familiars of the turnar (womb) found in over 20% of young worzes. There are firm, ball like and may vary in size from pin points to very large. Heedsds are diagnosed by the fisitory, gynecological examination and sompapiny. Their management depends upon their size and location, the ago of the patient, the symptoms the patient has and the patients desire for

There are Subsersous Birotis, which gove on the uper side of the uteria and intermual filestick, thick he in the middle of the worth. The Brods's thick he in the middle of the worth. The Brods's term of the subsersous term of the subsersous term. These grow on the inter side of the worth. When the best grows or the part which sheets during preprints. Does to mid subsersous Brods's can cause and the subsersous and difficulty in normal derivery. Most Brods to med surgey but for the first time in Most Brods to med surgey but for the first time in Most Brods to med surgey but for the first time in the subsersous and difficulty in normal derivery.

of filmends, without surgery
According to the Enhant Dallom Fal. commission
According to the Enhant Dallom Fal. Commission
According to the Enhant Dallom Fal.
Sept. Till now. Bleetide could cold be treated by
Engage and the Enhant Dallom Fall Color
Sept. Till now. Bleetide could cold be filmed
Compared in Municipal for the first time there be no
Enhant Dallom Fall Color
Enhant Dallom Fall Color
Enhant Dallom Fall
Enhant Dallom Fall
Enhant Dallom
Enhant

k the next day. formen in their 20's to those in their 50's, should be for nearly checkurs.

